

On ergodic behavior of p -adic dynamical systems ^{*}

Matthias Gundlach

Institut für Dynamische Systeme, Universität Bremen
Postfach 330 440, 28334 Bremen, Germany

Andrei Khrennikov[†] & Karl-Olof Lindahl [‡]

School of Mathematics and Systems Engineering
Växjö University, 351 95, Växjö, Sweden

Abstract

Monomial mappings, $x \mapsto x^n$, are topologically transitive and ergodic with respect to Haar measure on the unit circle in the complex plane. In this paper we obtain an analogous result for monomial dynamical systems over p -adic numbers. The process is, however, not straightforward. The result will depend on the natural number n . Moreover, in the p -adic case we never have ergodicity on the unit circle, but on the circles around the point 1.

1 Introduction

Investigations in p -adic quantum physics [14] – [23] (especially string theory [14], [15], [16]) stimulated an increasing interest in studying p -adic dynamical systems, see for example [1] – [12]. Some steps in this direction [23], [17] demonstrated that even the simplest (monomial) discrete dynamical systems over the fields of p -adic numbers $\mathbb{Q}_p$ have quite complex behavior. This behavior depends crucially on the prime number $p > 1$ (which determines $\mathbb{Q}_p$). By varying p we can transform attractors into centers of Siegel discs and vice versa. The number of cycles and their lengths also depend crucially on p [6], [13]. Other aspects (such as recurrence, renormalization and conjugacy) of monomials and the dynamics of the endomorphisms of the multiplicative group in general was investigated by Arrowsmith and Vivaldi [4].

Some applications of discrete p -adic dynamical systems to cognitive sciences and neural networks were considered in [23], [18]. Some of these cognitive models are described by random dynamical systems in the fields p -adic numbers, see [20], [24]. In this note we study ergodicity of monomial p -adic dynamical

^{*}This note was published in *Infinite Dimensional Analysis*, Vol. 4, No. 4 (2001) 569–577.

[†]This research was supported by the visiting professor fellowship at Bremen University.

[‡]This research was supported by the grant "Strategical investigations" of Växjö University.

systems on spheres. For a system $\psi_n(x) = x^n$, $n = 2, 3, \dots$, the result depends crucially on the relation between n and p . Our proof is essentially based on p -adic analysis (analytic mappings), [25]. We remark that the corresponding fact for the field of complex numbers, $\psi_n : \mathbb{C} \rightarrow \mathbb{C}$, for the sphere $|z| = 1$, is rather trivial, see [26].

2 p -adic numbers

The system of p -adic numbers $\mathbb{Q}_p$ was constructed by K. Hensel in the 1890s.

The field of real numbers $\mathbb{R}$ is constructed as the completion of the field of rational numbers $\mathbb{Q}$ with respect to the metric $\rho(x, y) = |x - y|$, where $|\cdot|$ is the usual valuation of distance given by the absolute value. The fields of p -adic numbers $\mathbb{Q}_p$ are constructed in a corresponding way, but using other valuations. For a prime number p , the p -adic valuation $|\cdot|_p$ is defined in the following way. First we define it for natural numbers. Every natural number n can be represented as the product of prime numbers, $n = 2^{r_2} 3^{r_3} \dots p^{r_p} \dots$, and we define $|n|_p = p^{-r_p}$, writing $|0|_p = 0$ and $|-n|_p = |n|_p$. We then extend the definition of the p -adic valuation $|\cdot|_p$ to all rational numbers by setting

$$|n/m|_p = |n|_p / |m|_p$$

for $m \neq 0$. The completion of $\mathbb{Q}_p$ with respect to the metric $\rho_p(x, y) = |x - y|_p$ is the locally compact field of p -adic numbers $\mathbb{Q}_p$.

The number fields $\mathbb{R}$ and $\mathbb{Q}_p$ are unique in a sense, since by Ostrovsky's theorem ([25]), $|\cdot|$ and $|\cdot|_p$ are the only possible valuations on $\mathbb{Q}$, but have quite distinctive properties. The field of real numbers $\mathbb{R}$ with its usual valuation satisfies $|n| = n \rightarrow \infty$ for valuations of natural numbers n and is said to be *Archimedean*. By a well know theorem of number theory [25], the only complete Archimedean fields are those of the real and the complex numbers. In contrast, the fields of p -adic numbers, which satisfy $|n|_p \leq 1$ for all $n \in \mathbb{N}$, are examples of *non-Archimedean* fields.

Unlike the absolute value distance $|\cdot|$, the p -adic valuation satisfies the strong triangle inequality

$$|x + y|_p \leq \max[|x|_p, |y|_p], \quad x, y \in \mathbb{Q}_p, \quad (1)$$

with equality in the case that $|x|_p \neq |y|_p$.

Write $B_r(a) = \{x \in \mathbb{Q}_p : |x - a|_p \leq r\}$ and $B_r^-(a) = \{x \in \mathbb{Q}_p : |x - a|_p < r\}$ where $r = p^n$ and $n = 0, \pm 1, \pm 2, \dots$. These are the “closed” and “open” balls in $\mathbb{Q}_p$ while the sets $S_r(a) = \{x \in K : |x - a|_p = r\}$ are the spheres in $\mathbb{Q}_p$ of such radii r . Any p -adic ball $B_r(0)$ is an additive subgroup of $\mathbb{Q}_p$, while the ball $B_1(0)$ is also a ring, which is called the *ring of p -adic integers* and is denoted by $\mathbb{Z}_p$.

As a consequence of the strong triangle property (1) every element b of a ball $B_r(a)$ is the center of this ball in the sense that $B_r(b) = B_r(a)$ for every $b \in B_r(a)$.

Each p -adic number has a unique expansion of the form $\sum_{n=M}^{\infty} a_n p^n$, where M is an integer and $a_n \in \{0, \dots, p-1\}$. The p -adic integers $\mathbb{Z}_p$, are p -adic numbers with expansions $\sum_{n=0}^{\infty} a_n p^n$. The absolute value of a p -adic number is p^{-k} if and only if k is the first digit in the expansion which differs from zero. Thus $|x - y|_p \leq p^{-k}$ if and only if $x \equiv y \pmod{p^k}$.

Let (m, n) denote the greatest common divisor of m and n . We say that n is a (multiplicative) *unit* (with respect to the prime number p) iff $(n, p) = 1$. Let G_{p^l} , $l \geq 1$, be the multiplicative group of units in the residue field modulo p^l . Let us by $\langle n \rangle = \{n^N : N \in \mathbb{N}\}$ denote the set *generated* by n .

Let ψ_n be a (monomial) mapping on $\mathbb{Z}_p$ taking x to x^n . Then all spheres $S_{p^{-l}}(1)$ are ψ_n -invariant iff n is a multiplicative unit. This is a consequence of the following result in p -adic analysis. A proof is given in Appendix A.

Lemma 2.1 *Let $x, y \in S_1(0)$ and suppose $|x - y|_p < 1$. Then for all natural numbers n ,*

$$|x^n - y^n|_p \leq |n|_p |x - y|_p, \quad (2)$$

with equality for $p > 2$. Moreover equality also holds for $p = 2$ if n is odd.

In particular, ψ_n is an isometry on $S_{p^{-l}}(1)$ if and only if $(n, p) = 1$. Therefore we will henceforth assume that n is a unit. Also note that, as a consequence, $S_{p^{-l}}(1)$ is not a group under multiplication. Thus our investigations are not about the dynamics on a compact (abelian) group.

3 Minimality

Let us consider the dynamical system $x \mapsto x^n$ on spheres $S_{p^{-l}}(1)$. The result depends crucially on the following well known result from group theory.

Lemma 3.1 *Let $p > 2$ and l be any natural number, then the natural number n is a generator of G_{p^l} if and only if n is a generator of G_{p^2} . G_{2^l} is noncyclic for $l \geq 3$.*

Recall that a dynamical system given by a continuous transformation ψ on a compact metric space X is called *topologically transitive* if there exists a dense orbit $\{\psi^n(x) : n \in \mathbb{N}\}$ in X , and (one-sided) *minimal*, if all orbits for ψ in X are dense. For the case of monomial systems $x \mapsto x^n$ on spheres $S_{p^{-l}}(1)$ topological transitivity means the existens of an $x \in S_{p^{-l}}(1)$ s.t. each $y \in S_{p^{-l}}(1)$ is a limit point in the orbit of x , i.e. can be represented as

$$y = \lim_{k \rightarrow \infty} x^{n^{N_k}}, \quad (3)$$

for some sequence $\{N_k\}$, while minimality means that such a property holds for any $x \in S_{p^{-l}}(1)$. Our investigations are based on the following theorem.

Theorem 3.1 *For $p \neq 2$ the set $\langle n \rangle$ is dense in $S_1(0)$ if and only if n is a generator of G_{p^2} .*

Proof. We have to show that for every $\epsilon > 0$ and every $x \in S_1(0)$ there is a $y \in \langle n \rangle$ such that $|x - y|_p < \epsilon$ if and only if n is a generator of G_{p^2} . First, assume that n is a generator of G_{p^2} . Let $\epsilon > 0$ and $x \in S_1(0)$ be arbitrary. Because of the discreteness of the p -adic metric we can assume that $\epsilon = p^{-k}$ for some natural number k . But (according to Lemma 3.1) if n is a generator of G_{p^2} , then n is also a generator of G_{p^l} for every natural number l (and $p \neq 2$) and especially for $l = k$. Consequently there is an N such that $n^N = x \pmod{p^k}$. From the definition of the p -adic metric we have that $|x - y|_p < p^{-k}$ if and only if $x \equiv y \pmod{p^k}$. Hence $|x - n^N|_p < p^{-k}$. By reversing the arguments above, we see that the converse is also true. $\square$

Let us consider $p \neq 2$ and for $x \in B_{p^{-1}}(1)$ the p -adic exponential function $t \mapsto x^t$, see, for example [25]. This function is well defined and continuous as a map from $\mathbb{Z}_p$ to $\mathbb{Z}_p$. In particular, for each $a \in \mathbb{Z}_p$, we have

$$x^a = \lim_{k \rightarrow a} x^k, \quad k \in \mathbb{N}. \quad (4)$$

We shall also use properties of the p -adic logarithmic function, see, for example [25]. Let $z \in B_{p^{-1}}(1)$. Then $\log z$ is well defined. For $z = 1 + \lambda$ with $|\lambda|_p \leq 1/p$, we have:

$$\log z = \sum_{k=1}^{\infty} \frac{(-1)^{k+1} \Delta^k}{k} = \lambda(1 + \lambda \Delta_\lambda), \quad |\Delta_\lambda|_p \leq 1. \quad (5)$$

By using (5) we obtain that $\log : B_{p^{-1}}(1) \rightarrow B_{p^{-1}}(0)$ is an isometry:

$$|\log x_1 - \log x_2|_p = |x_1 - x_2|_p, \quad x_1, x_2 \in B_{1/p}(1). \quad (6)$$

Lemma 3.2 *Let $x \in B_{p^{-1}}(1), x \neq 1, a \in \mathbb{Z}_p$ and let $\{m_k\}$ be a sequence of natural numbers. If $x^{m_k} \rightarrow x^a, k \rightarrow \infty$, then $m_k \rightarrow a$ as $k \rightarrow \infty$, in $\mathbb{Z}_p$.*

This is a consequence of the isometric property of $\log$.

Theorem 3.2 *Let $p \neq 2$ and $l \geq 1$. Then, the monomial dynamical system $x \mapsto x^n$ is minimal on the circle $S_{p^{-l}}(1)$ if and only if n is a generator of G_{p^2} .*

Proof. Let $x \in S_{p^{-l}}(1)$. Consider the equation $x^a = y$. We have that $a = \frac{\log x}{\log y}$. As $\log : B_{p^{-1}}(1) \rightarrow B_{p^{-1}}(0)$ is an isometry, we have $\log(S_{p^{-l}}(1)) = S_{p^{-l}}(0)$. Thus $a = \frac{\log x}{\log y} \in S_1(0)$ and moreover, each $a \in S_1(0)$ can be represented as $\frac{\log x}{\log y}$ for some $y \in S_{p^{-l}}(1)$.

Let y be an arbitrary element of $S_{p^{-l}}(1)$ and let $x^a = y$ for some $a \in S_1(0)$. By Theorem 3.1, if n is a generator of G_{p^2} , then each $a \in S_1(0)$ is a limit point of the sequence $\{n^N\}_{N=1}^{\infty}$. Thus $a = \lim_{k \rightarrow \infty} n^{N_k}$ for some subsequence $\{N_k\}$. By using the continuity of the exponential function we obtain (3).

Suppose now that, for some n , $x^{n^{N_k}} \rightarrow x^a$. By Lemma 3.2 we obtain that $n^{N_k} \rightarrow a$ as $k \rightarrow \infty$. If we have (3) for all $y \in S_{p^{-l}}(1)$, then each $a \in S_1(0)$ can be approximated by elements n^N . In particular, all elements in the set $\{1, 2, \dots, p-1, p+1, \dots, p^2-1\}$ can be approximated modulo p^2 . Thus n is a generator of G_{p^2} . $\square$.

Example In the case that $p = 3$ we have that ψ_n is minimal if $n = 2$, 2 is a generator of $U_{3^2} = \{1, 2, 4, 5, 7, 8\}$. But for $n = 4$ it is not; $\langle 4 \rangle \bmod 3^2 = \{1, 4, 7\}$. We can also see this by noting that $S_{1/3}(1) = B_{1/9}(4) \cup B_{1/9}(7)$ and that $B_{1/9}(4)$ is invariant under ψ_4 .

Corollary 3.1 *If a is a fixed point of the monomial dynamical system $x \mapsto x^n$, then this is minimal on $S_{p^{-l}}(a)$ if and only if n is a generator of G_{p^2} .*

Proof. The assertion follows immediately from Theorem 3.2 by topological conjugation via $x \mapsto x/a$, $S_{p^{-l}}(a) \mapsto S_{p^{-l}}(1)$. $\square$

Example Let $p = 17$ and $n = 3$. On $\mathbb{Q}_{17}$ there is a primitive 3rd root of unity, see for example [24]. Moreover, 3 is also a generator of G_{17^2} . Therefore there exist n th roots of unity different from 1 around which the dynamics is minimal.

4 Unique ergodicity

In the following we will show that the minimality of the monomial dynamical system $\psi_n : x \mapsto x^n$ on the sphere $S_{p^{-l}}(1)$ is equivalent to its *unique ergodicity*. The latter property means that there exists a unique probability measure on $S_{p^{-l}}(1)$ and its Borel σ -algebra which is invariant under ψ_n . We will see that this measure is in fact the normalized restriction of the Haar measure on $\mathbb{Z}_p$. Moreover, we will also see that the ergodicity with respect to Haar measure of ψ_n is also equivalent to its unique ergodicity. We should point out that – though many results are analogous to the case of the (irrational) rotation on the circle, our situation is quite different, in particular as we do not deal with dynamics on topological subgroups.

Lemma 4.1 *Assume that ψ_n is minimal. Then, the Haar measure m is the unique ψ_n -invariant measure on $S_{p^{-l}}(1)$.*

Proof. First note that minimality of ψ_n implies that $(n, p) = 1$ and hence that ψ_n is an isometry on $S_{p^{-l}}(1)$, see equation (2). Then, as a consequence of Theorem 27.5 in [25], it follows that $\psi_n(B_r(a)) = B_r(\psi_n(a))$ for each ball $B_r(a) \subset S_{p^{-l}}(1)$. Consequently, for every open set $U \neq \emptyset$ we must have that $S_{p^{-l}}(1) = \bigcup_{N=0}^{\infty} \psi_n^N(U)$. Then, it follows for a ψ_n -invariant measure μ that $\mu(U) > 0$.

Moreover, we can split $S_{p^{-l}}(1)$ into disjoint balls of radii $p^{-(l+k)}$, $k \geq 1$, on which ψ_n acts as a permutation. In fact, for each $k \geq 1$, $S_{p^{-l}}(1)$ is the union,

$$S_{p^{-l}}(1) = \bigcup B_{p^{-(l+k)}}(1 + b_l p^l + \dots + b_{l+k-1} p^{l+k-1}), \quad (7)$$

where $b_i \in \{0, 1, \dots, p-1\}$ and $b_l \neq 0$.

We now show that ψ_n is a (transitive) permutation on the partition (7). Recall that every element of a p -adic ball is the center of that ball (see section 2), and as pointed out above $\psi_n(B_r(a)) = B_r(\psi_n(a))$. Consequently we have for all positive integers k , $\psi_n^k(a) \in B_r(a) \Rightarrow \psi_n^k(B_r(a)) = B_r(\psi_n^k(a)) = B_r(a)$ so that $\psi_n^{Nk}(a) \in B_r(a)$ for every natural number N . Hence, for a minimal ψ_n a point of a ball B of the partition (7) must move to another ball in the partition. Furthermore, the minimality of ψ_n shows indeed that ψ_n acts as a permutation on balls. By invariance of μ , all balls must have the same positive measure. As this holds for any k , μ must be the restriction of Haar measure m . $\square$

The arguments of the proof of Lemma 4.1 also show that Haar measure is always ψ_n -invariant. Thus if ψ_n is uniquely ergodic, the unique invariant measure must be the Haar measure m . Under these circumstances it is known ([26, Theorem 6.17]) that ψ_n must be minimal.

Theorem 4.1 *The monomial dynamical system $\psi_n : x \mapsto x^n$ on $S_{p^{-l}}(1)$ is minimal if and only if it is uniquely ergodic in which case the unique invariant measure is the Haar measure.*

Let us mention that unique ergodicity yields in particular the ergodicity of the unique invariant measure, i.e. the Haar measure m , which means that

$$\frac{1}{N} \sum_{i=0}^{N-1} f(x^{n^i}) \rightarrow \int f dm \text{ for all } x \in S_{p^{-l}}(1), \quad (8)$$

and all continuous functions $f : S_{p^{-l}}(1) \rightarrow \mathbb{R}$.

On the other hand the arguments of the proof of Lemma 4.1, i.e. the fact that ψ_n acts as a transitive permutation on each partition of $S_{p^{-l}}(1)$ into disjoint balls if and only if $\langle n \rangle = G_{p^2}$, proves that if n is not a generator of G_{p^2} then the system is not ergodic with respect to Haar measure. Consequently, if ψ_n is ergodic then $\langle n \rangle = G_{p^2}$ so that the system is minimal by Theorem 3.2, and hence even uniquely ergodic by Theorem 4.1. Since unique ergodicity implies ergodicity one has the following.

Theorem 4.2 *The monomial dynamical system $\psi_n : x \mapsto x^n$ on $S_{p^{-l}}(1)$ is ergodic with respect to Haar measure if and only if it is uniquely ergodic.*

Even if the monomial dynamical system $\psi_n : x \mapsto x^n$ on $S_{p^{-l}}(1)$ is ergodic, it cannot be mixing, especially not weak-mixing. This can be seen from the fact that an abstract dynamical system is weak-mixing if and only if the product of such two systems is ergodic. If we choose a function f on $S_{p^{-l}}(1)$ and define a function F on $S_{p^{-l}}(1) \times S_{p^{-l}}(1)$ by $F(x, y) := f(\log x / \log y)$ (which is well defined as $\log$ does not vanish on $S_{p^{-l}}(1)$), we obtain a non-constant function satisfying $F(\psi_n(x), \psi_n(y)) = F(x, y)$. This shows (see [26, Theorem 1.6]) that $\psi_n \times \psi_n$ is not ergodic, and hence ψ_n is not weak-mixing with respect to any invariant measure, in particular not to the Haar measure.

Let us consider the ergodicity of a perturbed system

$$\psi_q = x^n + q(x), \quad (9)$$

for some polynomial q such that $q(x) \equiv 0 \pmod{p^{l+2}}$, $(|q(x)|_p \leq p^{-(l+2)})$. Under this condition, the sphere $S_{p^{-l}}(1)$ is invariant. For such a system to be ergodic it is necessary that n is a generator of G_{p^2} . This follows from the fact that for each $x = 1 + a_l p^l + a_{l+1} p^{l+1} + \dots$ on $S_{p^{-l}}(1)$ the condition on q gives

$$\psi_q^N(x) \equiv 1 + n^N(a_l + a_{l+1}p)p^l \pmod{p^{l+2}}. \quad (10)$$

Now, ψ_q acts as a transitive permutation on the $p(p-1)$ balls of radius $p^{-(l+2)}$ if and only if $\langle n \rangle = G_{p^2}$. Consequently, a perturbation (9) cannot make a nonergodic system ergodic. However, it is not clear at this point if this is true also if $|q(x)| = p^{-(l+1)}$.

5 Acknowledgement

We would like to thank P.A. Svensson at Växjö Univerity for consultation in algebra.

A Appendix

Lemma A.1¹ *Let $x, y \in S_1(0)$ and suppose $|x - y|_p < 1$. Then for all natural numbers n ,*

$$|x^n - y^n|_p \leq |n|_p |x - y|_p, \quad (11)$$

with equality for $p > 2$. Moreover equality also holds for $p = 2$ if n is odd.

Proof. First note that

$$\begin{aligned} |x^n - y^n|_p &= |(y + x - y)^n - y^n|_p = \left| \sum_{k=0}^n \binom{n}{k} y^{n-k} (x - y)^k - y^n \right|_p \\ &= \left| \sum_{k=1}^n \binom{n}{k} y^{n-k} (x - y)^k \right|_p \leq \max_k \left| \binom{n}{k} y^{n-k} (x - y)^k \right|_p, \end{aligned}$$

where the inequality is a consequence of the strong triangle inequality (1). We will see that the maximum occurs for $k = 1$ and that the corresponding term is greater than every other term (except for the case where $p = 2$ and n is even). By the multiplicative property of the valuation, the valuation of the k th term

¹This result was obtained in [24].

in the sum can be estimated according to

$$\begin{aligned}
\left| \binom{n}{k} y^{n-k} (x-y)^k \right|_p &= \left| \binom{n}{k} \right|_p |y|_p^{n-k} |x-y|_p^k \\
&= |n(n-1) \cdots (n-k+1)|_p |x-y|_p \frac{|x-y|_p^{k-1}}{|k!|_p} \\
&\leq |n|_p |x-y|_p \frac{|x-y|_p^{k-1}}{|k!|_p} \\
&= \left| \binom{n}{1} y^{n-1} (x-y)^1 \right|_p \frac{|x-y|_p^{k-1}}{|k!|_p},
\end{aligned}$$

where the inequality holds because $|a|_p \leq 1$ for every integer $a \in \mathbb{Z}$ (together with the multiplicative property of the valuation). Note that strong inequality holds in the case that $p = 2$ and n is odd; if n is odd we have $|n|_2 = 1$ (n is not divisible by 2). Moreover we can assume that $n > 1$ (the lemma is automatically true for $n = 1$) so that $|n-1|_2 < 1$.

To estimate the latter expression, $|x-y|_p^{k-1} / |k!|_p$, we use the fact that $|k!|_p \geq 1/p^{k-1}$ with strong inequality for $p > 2$, see for example the paper [25, Lemma 25.5]. Moreover $|x-y|_p \leq 1/p$ by the condition of the lemma. Consequently,

$$|x-y|_p^{k-1} \leq 1/p^{k-1}.$$

Hence we have that $|x-y|_p^{k-1} / |k!|_p \leq 1$ with strong inequality when $p > 2$. Therefore, for every integer k such that $1 \leq k \leq n$, we have

$$\left| \binom{n}{k} y^{n-k} (x-y)^k \right|_p \leq \left| \binom{n}{1} y^{n-1} (x-y)^1 \right|_p,$$

with strong inequality for all p and n except the case when $p = 2$ and n is even. Hence, as a consequence of the strong triangle inequality (1) we have that

$$|x^n - y^n|_p \leq \left| \binom{n}{1} y^{n-1} (x-y)^1 \right|_p = |n|_p |x-y|_p,$$

with equality for $p > 2$ and, if n is odd, also for $p = 2$. $\square$

References

- [1] Thiran E., Verstegen D., Weyers J., p -adic dynamics, *J. Stat. Phys.* **54** (1989), 893-913.
- [2] Ben-Menahem, S., p -adic iterations (Tel Aviv Univ. Press, Tel Aviv, 1988), preprint TAUP 1627-88.

- [3] Arrowsmith, D.K. & Vivaldi, F., Some p -adic representations of the Smale horseshoe, *Physics letters A* **176** (1993), 292-294.
- [4] Arrowsmith, D.K. & Vivaldi, F., Geometry of p -adic Siegel discs, *Physica D* **74** (1994), 222-236.
- [5] Lubin, J., Non-Archimedean dynamical systems, *Compositio Mathematica*, **94**, 1994, 321-346.
- [6] Pezeda, T., Polynomial cycles in certain local domains, *Acta Arithmetica*, LXVI, 1994, 11-22.
- [7] Lubin, J., Sen's theorem on iteration of power series, *Proc. Am. Math Soc.*, **123**, 1995, 66-99.
- [8] Woodcock, F., Smart, P., p -adic chaos and random number generation, *Experimental Mathematics*, **7**, 1998, 334-342.
- [9] Benedetto, R., Fatou components in p -adic dynamics, PHD thesis, Brown University, 1998.
- [10] Benedetto, R., Hyperbolic maps in p -adic dynamics, Preprint, University of Rochester, 1999.
- [11] Hsia, L., A weak Néron model with applications to p -adic dynamical systems, *Compositio Mathematica*, **100**, 1996, 227-304.
- [12] Bosio, D. & Vivaldi, F., Round-off errors and p -adic numbers, *Nonlinearity*, **13**, 2000, 309-322.
- [13] Nilsson, M., Cycles of monomial dynamical systems over the field of p -adic numbers, Reports from Växjö University, **20**, 1999.
- [14] Volovich I. V., p -adic string. *Class. Quant. Grav.*, **4**, 83-87 (1987).
- [15] Freund P. G. O. and Witten E., Adelic string amplitudes. *Phys. Lett. B*, **199**, 191-195 (1987).
- [16] Vladimirov V. S., Volovich I. V., and Zelenov E. I., *p -adic Analysis and Mathematical Physics*, World Scientific Publ., Singapore (1994).
- [17] Albeverio S., Khrennikov, A. Yu., De Smedt, S., Tirozzi, B., p -adic dynamical systems, *Theor. and Math. Phys.*, **114**, N. 3, 1998, 3-20.
- [18] Albeverio, S., Khrennikov, A. Yu., Kloeden, P. Memory retrieval as a p -adic dynamical system. *Biosystems*, **49**, 105-115, (1999).
- [19] Albeverio S., Khrennikov, A. Yu., Tirozzi, B., p -adic neural networks. To be published.

- [20] Dubischar D., Gundlach, V.M., Steinkamp, O., Khrennikov, A. Yu., Attractors of random dynamical systems over p -adic numbers and a model of noisy cognitive processes. *Physica D*, **130**, 1-12, (1999).
- [21] Gundlach, V.M., Khrennikov, A., Lindahl, K.O., Ergodicity on p -adic sphere, German Open Conference on Probability and Statistics, University of Hamburg, March 21-24, 2000, p.61.
- [22] Khrennikov A., *p -adic Valued Distributions in Mathematical Physics*. Kluwer, 1994.
- [23] Khrennikov A., *Non-Archimedean Analysis: Quantum Paradoxes, Dynamical Systems and Biological Models*. Kluwer, 1997.
- [24] Lindahl K.O., On Markovian properties of the dynamics on attractors of random dynamical systems over p -adic numbers. Reports from MASDA, Växjö University, June 1999.
- [25] W.H. Schikhof, *Ultrametric calculus*, Cambridge University press, 1984.
- [26] Walters, P. *An introduction to ergodic theory*. Springer, 1982.